



ORANGE COUNTY EMERGENCY MEDICAL SERVICES
BASE HOSPITAL TREATMENT GUIDELINES
ADULT/ADOLESCENT

BH-P-010
Page: 1 of 2
Org. Date: 4/01/2017
Revise Date: 10/01/2017

NEWBORN IN THE FIELD

BASE GUIDELINES	ALS STANDING ORDER
<p>1. Hospitals with Obstetrical Service (appropriate for triage of mother and newborn):</p> <p>Anaheim Global Medical Center Anaheim Regional Medical Center Fountain Valley Regional Hospital Garden Grove Medical Center Hoag Memorial Hospital Kaiser (Anaheim) Kaiser (Irvine) La Palma Intercommunity Hospital Los Alamitos Medical Center Mission Hospital, Mission Viejo Orange Coast Memorial Medical Center Orange County Global Medical Center Saddleback Memorial Medical Center St. Joseph Hospital St. Jude Medical Center South Coast Global Medical Center UCI Medical Center</p> <p>2. Once cord is clamped, it does not necessarily need to be cut in the field.</p> <p>3. If a nuchal cord occurs upon delivery, instruct the paramedic to slip a finger under the cord and move it over the head to free the cord, slight to moderate stretching the cord to accomplish this maneuver is appropriate.</p>	<p>1. If possible, deliver infant in a warm, draft-free area.</p> <p>2. Minimize heat loss: dry face, head, and body with soft towel or blanket. Do not allow wet linen to remain in contact with child. Wrap infant in dry blanket or towel.</p> <p>3. Position infant on side with neck in a neutral position to allow secretions to collect in the mouth to drain rather than accumulate the posterior pharynx.</p> <p>4. Assure airway is open; provide oxygen by blow-by technique until child is active and crying.</p> <p>5. Perform Apgar Score at 1 minute and 5 minutes after birth (see Guidelines below).</p> <p>6. If possible, clamp and cut cord (see OCEMS Procedure B-060).</p> <p>7. Contact Base Hospital for appropriate destination with neonatal care capability.</p> <p>8. ALS escort infant and mother.</p> <p>NEWBORN IN DISTRESS:</p> <p><u>If respiratory depression:</u></p> <p>A. Assure airway is open and clear of secretions.</p> <p>B. Administer oxygen by blow-by technique.</p> <p>C. Stimulate by rubbing along the spine or soles of feet.</p> <p>D. Reposition to assure open airway.</p> <p>E. For respiratory depression due to suspected maternal use of narcotics and unresponsive to stimulation: ▶ <i>Naloxone 0.1 mg/kg IN/IM once to a maximum of 1 mg.</i></p> <p>F. Contact Base Hospital for appropriate destination with neonatal care capability.</p> <p><u>Gasping Respirations / Apnea / HR<100/minute / Persistent Cyanosis:</u></p> <p>A. Assisted positive pressure ventilation with appropriate sized bag-valve-mask, rate 40-60 breaths/minute with 100% oxygen.</p> <p>B. Contact Base Hospital for appropriate destination with neonatal care capability.</p>

Approved: 



**ORANGE COUNTY EMERGENCY MEDICAL SERVICES
BASE HOSPITAL TREATMENT GUIDELINES
ADULT/ADOLESCENT**

BH-P-010
Page: 2 of 2
Org. Date: 4/01/2017
Revise Date: 10/01/2017

NEWBORN IN THE FIELD

BASE GUIDELINES

APGAR SCORE

- The Apgar Score measures newborn's status.
- Each of the five categories are scored from 0-2 and then totaled. Apgar scoring is done twice; 1 minute after birth and 5 minutes after birth.
- Resuscitation, if needed, should not await Apgar scoring.

Scoring	0	1	2
Heart rate	Absent	Slow (below 100)	≥ 100
Respiratory Effort	Absent	Weak Cry; Hypoventilation	Strong Cry
Muscle Tone	Limp	Slight Flexion of Extremities	Active Flexion
Reflex Responses (to bulb syringe in nostril)	No response	Some Grimace	Cough or cry
Color	Blue, pale	Body pink, Extremities blue	Completely pink

- Total score indications are:
 - One (1) minute Apgar Score indications:
 - 7-10 – a healthy infant.
 - 4-6 – a potentially sick infant.
 - 0-3 – a severely depressed infant.
 - Five (5) minute Apgar Score indications:
 - 7-10 – a healthy infant.
 - 4-6 – a potentially sick infant.
 - 0-3 – a severely depressed infant.

ALS STANDING ORDER

Heart Rate < 60/minute / No Response to Above Measures:

- Begin chest compressions; interpose chest compressions with ventilations in a 3:1 ratio (90 compressions and 30 breaths per minute).
- Contact Base Hospital for appropriate destination with neonatal care capability.

TREATMENT GUIDELINES:

- Refer to OCEMS Procedure B-060 for steps in field delivery of an infant.
- Drying will usually produce enough stimulation to initiate effective respirations in most newborns.
- To determine heart rate: auscultate, do not attempt to palpate.

Approved:

Reviewed:9/2017
Final Date for Implementation: Oct. 1, 2017
OCEMS copyright © 2017